















- [8] Hoogeveen RC, Gaubatz JW, Sun W, Dodge RC, Crosby JR & Jiang J. (2014): Small Dense Low-Density Lipoprotein-Cholesterol Concentration Predict Risk for Coronary Heart Disease Significance. *Arterioscler Thromb. Vasc. Biol.* 34:1-9.
- [9] Lahiri DK & Nurnberger JI. (1991): A rapid non-enzymatic method for the preparation of HMW DNA from blood for RFLP studies. *Nucleic Acids Res.* 19: 5444.
- [10] Lin N, Lee M, Lee R, Hong AIC & Chen HI. (2008): Analysis of Endothelial Nitric Oxide Synthase Gene Polymorphisms with Cardiovascular Diseases in Eastern Taiwan. *Chin J Physiol.* 51:42-47.
- [11] Litwack G. (2014): Nitric Oxide. Elsevier Science . 96 : 390- 394.
- [12] Mahmoodi K, Nasehi L, Karami E & Soltanpour MS. (2016): Association of Nitric Oxide Levels and Endothelial Nitric Oxide Synthase G894T Polymorphism with Coronary Artery Disease in the Iranian Population. *ncbi.nlm.nih.gov.* 32: 105-112.
- [13] Marsden PA, Heng HH, Scherer SW, Sterwart RJ, Hall A V & Shi XM. (1993): Structure and chromosomal of the human constitution endothelial nitric oxide synthase gene. *J.Biol. Chem.* 268: 17478-17488.
- [14] Mayer B & Hemmens B. (1997): Biosynthesis and action of nitric oxide in mammalian cells. *Trends Biochem. Sci.* 22:477-481.
- [15] Meroufel D, Mediene- benchekor S & Dumont J. (2009): Relationship between Endothelial nitric oxide synthase gene polymorphisms and the risk of myocardial Infraction in the Algerian population. *Egy. Jou. Med. H. Gen.* 10:89-95.
- [16] Nakayama M, Yasue H, Yoshimura M, Shimasaki Y, Kugiyama K & Ogawa H. (1999): T -786→C Mutation in the 5-Flanking Region of the Endothelial Nitric Oxide Synthase Gene Is Association With Coronary Spasm. *Circulation.* 99:2846-2870.
- [17] Oemar BS, Tschudi MR, Gooy N, Brovkovich V, Malinski T & Liischer TE. (1998): Reduced endothelial nitric oxide synthase expression and production in human atherosclerosis. *Circulation.* 30: 2494-2498.
- [18] Prabhakar P, Thatte HS, Goetz RM, Cho MR, Golan DE & Michel T. (1998): Receptor -regulated translocation of endothelial nitric-oxide synthase. *J. Biol. Chem.* 16:27383- 27388.
- [19] Puddu P, Cravero E, Puddu GM & Muscari. (2005): A Genes and atherosclerosis at The region of the predisposition. *Int.J. Clin. Pract.* 59:462-472.
- [20] Salazar LA, Hirata MH, Forti N, Diament Jm Giannini SD, Issa JS & Hirata RD. (2000): Polymorphism at the LDL receptor gene is associated with differences in serum lipid concentration in subjects with low and high risk for coronary artery disease from Brazil. *Clin. Chim. Acta.* 293:75-88.
- [21] Salimi S, Naghavi A, Firoozrai M, Zand H, Trvilani H & Nakhaee A. (2012): Association of plasma nitric oxide concentration and endothelial nitric oxide synthase T-786C gene polymorphism in coronary artery diseas *Pathophysiology* 19:157-162.
- [22] Searles CD. (2006): Transcriptional and posttranscriptional regulation of endothelial nitric oxide synthase expression. *Am J Physiol – Cell Physiol.* 291:803-816.
- [23] Sigusch HH, Surber R, Lehmann MH, Surber S, Weber J & Henke A. (2000): Lack of association between 27-bp repeat polymorphism in intron 4 of the endothelial nitric oxide synthase gene and the risk of coronary artery disease. *Scand. J. Clin. Lab. Investig.* 60:229-235.
- [24] Vanin AF.(1998): Biological role of nitric oxide: history, modern state, and perspectives for research. *Bio. M. J.* 63: 731-733.
- [25] Wang X, Sim AS, Badenhop RF, Mccerdie RM & Wilcken DE. (1996): A smoking-dependent risk of coronary artery disease associated with a polymorphism of the endothelial nitric oxide synthase gene. *Nat. Med.* 2:41-45.